


--	--	--	--	--

[illegible]

1961 TIME Ancel Keys

[illegible]



1984年2月13日  
 第251卷第3期

Lipid代谢与“饮食与健康”

JAMA, Jan 20, 1984 - Vol 251, No.3

1

# The Lipid Research Clinics Coronary Primary Prevention Trial Results

## I. Reduction in Incidence of Coronary Heart Disease

## Lipid Research Clinics Program

[illegible][illegible]

death and/or definite nonfatal myocardial infarction—reflecting a 24% reduction in definite CHD death and a 19% reduction in nonfatal myocardial infarction. The cumulative seven-year incidence of the primary end point was 7% in the cholestyramine group v 8.6% in the placebo group. In addition, the incidence rates for new positive exercise tests, angina, and coronary bypass surgery were reduced by 25%, 20%, and 21%, respectively, in the cholestyramine group. The risk of death from all causes was only slightly and not significantly reduced in the cholestyramine group. The magnitude of this

--	--	--	--	--	--

3806

Lipid 

[illegible][illegible]Lipid 

--	--	--	--	--	--	--	--

24%☐☒☐☐☐☐

[illegible]

24%

24%


--	--	--	--	--

[illegible]

24%

Lipid 

	LDL	HDL
LDL	1900	1906
LDL	216.2	215.6
7mmol/L	197.6	174.9
mmol/L	38	30
mmol/L	2%	1.57%
mmol/L	71	68
mmol/L	3.73%	3.57%



24%  $\frac{0.4\%}{1.6\%} = 24\%$




24%  $\frac{0.4\%}{1.6\%} = 24\%$







Ox-LDL structure: 28 units (Apolipoprotein B100) + 5 units (Hydrophobic core) = Ox-LDL



.

Revision #2  
Created 1 August 2021 12:17:48 by [ ]  
Updated 1 August 2021 12:24:28 by [ ]